Analysis Sandalwood, Royal Hawaiian Essential Oil

Laboratoire PhytoChemia









Date: June 14, 2017

SAMPLE IDENTIFICATION

Internal code: 17F08-FSE3-1-DM

Customer Identification: Sandalwood, Royal Hawaiian - Lot #HA-1132016

Type: Essential oil

Source: Santalum paniculatum

Customer: Flowers Shining Everywhere Inc.

livinglibation

ANALYSIS

Method: PC-PA-001-15E06, "Analysis of the composition of a liquid essential oil by GC-FID" (in French).

Analyst: Sylvain Mercier, M. Sc., chimiste

Analysis date: 2017-06-13

Checked and approved by:

livinglibgtign

Alexis St-Gelais, M. Sc., chimiste 2013-174

livinglibation/

Note: This report may not be published, including online, without the written consent from Eaboratoire PhytoChemia.

This report is digitally signed, it is only considered valid if the digital signature is intact.

livinglibotion

livinglibation

livinglibation

ivinglibotion

livinglibation



IDENTIFIED COMPOUNDS

44.0	200			
IIVI	ngi	100	uo	TH.

livingilbottoru	Column: BP5			Column: WAX				
Identification	R.T.	R.L	96	90	RJ.	R.T.	Molecular Class	
Santene	2.62	876	0.28	0.26	881	0.84	Normonoterpene	
Tricyclene	3.22	917	0.01	0.01	912	0.95	Monoterpene	
Teresantalic acid	13.70	1300	0.38	0.45	2128	39.78	Monoterp, acid	
Tricycloekasantalal	15.85	1333	0.15				Terpenic aldehyde	
a-Cedrene	20.11	1398	0.11	0.07	1481	8.19	Sesquiterpene	
a-Funebrene liv	2039	CT401	0.02	0.02	1511	8.90	Sesquiterpene	
g-Santalene	20.87	1407	0.71	0.70	1502	8.69	Sesquiterpene	
trans-a-Bergamotene	22.15	1423	0.13	0.10	1521	9.13	Sesquiterpene	
(Z)-epl-β-Santalene	23.08	1434	0.60	0.52	1556	10.36	Sesquiturpene	
(Z)-B-Santalene	24.10	1446	0.93	0.96	1571	10.88	Sesquiterpena	
y-Curcumene	25.89	1468	0.17	0.14	1616	12.54	Sesquiterpene	
ar Curcumene	26.40	1474	0.17	0.20	1687	16.01	Sesquiterpene	
9o-Seinene	26.71	1477	0.11	0.07	1635	13.34	Sesquiterpene	
- B-Bisabolene	28.70	1502	0.29	0.25	1652	14.21	Sesquiterpene	
β-Curcumene	28.84	1504	0.32	0.24	1662	14.75	Sesquiterpene	
(E)-Nerolidol	33.67	1571	0.13	0.21	1988	34.65	Sesquiterp, alcohol	
Dendrolasin	33,77	100 mg	0.00	0.14	1853	26.30	Sesquiterp, ether	
g-Acorenol	36.31	1613	0.21	0.15	2020	36.10	Sesquiterp, alcohol	
B-Acorenol	36.64	1621	0.11	0.11	2042	37,03	Sesquiterp, alcohol	
Unknown (m/z = 91, 179	37.42	1640	0.53	10.000.00.00	1,50,510,51	30.000	Oxygenated sesquiterp.	
(62), 79 (58), 105 (58), 93								
(58), 67 (57), 161 (57) 188						living	alibalianz	
(42), 216 (6))								
g-Bisabolol oxide B	37.59	1645	0.47	0.47	2034	36.75	Sesquiterp, alcohol	
Unknown (m/z = 95, 121	37.64	1646	0.25	n/Joseph	-		Oxygenated sesquiterp.	
(84), 161 (77), 43 (63), 79 (61), 91 (61), 105 (54), 93								
(54), 294 (47) 220 (8)) (E)-q-Santalai?	20.10	11450					Committee of the Control	
	38.18	1659	0.94	tions:			Sesquiterp. aldehyde	
Santalal analog?	38.28	1000000		The same of	and the second		Sesquiterp, aldehyde	
Cedr-8-en-13-ol	38.70	1672	+ 4.02	3.56	2258	43.19	Sesquiterp, alcohol	
(Z)-o-Santalol	39.11	1682	28.23	28.94	2249	42.96	Sesquiterp. alcohol #	
(Z)-a-trans-Bergamotol	39.54*	1692	3.98	0.95	2277	43.62	Sesquiterp, alcohol	
(£)-o-Santalol	39.54"	1692	[3.98]	2.48	2306	44.27	Sesquiterp, alcohol	
(Z)-epi-β-Santalol	39.99	1704	2.78	11.55	2325	44.69*	Sesquiterp. alcohol	
(Z) (II-Santal@Lottion/	40.55	1722	10.93	[11.55]	2325	44.69*		
(Z)-Nuciferol	40.79	1730	4.89	8.65	2419	46.71	Sesquiterp. alcohol	
(Z)-y-Curcumen-12-ol	40.94	1734	6.52	3.76	2342	45.06	Sesquiterp. alcohol	
(£)-β-Santalol	41.29	3845n	9 573	tions		1000000	Sesquiterp. alcohol	
(Z)-β-Curcumen-12-ol	41.86	1763	4.78	5.00	2389	46.09	Sesquiterp, alcohol	
(Z)-Lanceol	42.13	1772	7.64	7.27	2397	46.26	Sesquiterp alcohol	

Laboratoire





(E)-Lanceol?	42.97	1798	0.48	0.45	2432	47.00	Sesquiterp. alcohol
Total identified			82.97%	77.68%	maye	NAME.	1D

^{*:} Two or more compounds are coeluting on this column

[xx]: Duplicate percentage due to coelutions, not taken account in the identified total

Note: no correction factor was applied

OTHER DATA

Physical aspect: Light yellow liquid

Refractive index: 1.5063 ± 0.0903 (20°C)

CONCLUSION

No adulterant, contaminant or diluent were detected using this method. The concentrations of santalois reported in literature are however slightly higher for *S. paniculatum*, with (2)-a-santaloi reported minimally at 34%. This oil however is not widely described in literature, and the values observed here could still be within natural variation.

REFERENCES

 Braun, N. A.; Sim, S.; Kohlenberg, B.; Lawrence, B. M. Hawaiian Sandalwood: Oil Composition of Santalum Paniculatum and Comparison with Other Sandal Species. Nat. Prod. Commun. 2014, 9 (9), 1365–1368.

livinglibation

livinglibation

livinglibation


livinglibations

livinglibation

livinglibations

Laboratoire PhytoChemia livinglibations

Plus que des analyses... des conseils



Plus que des analyses... des conseils

Page 4 of 5

